

PRIOR AUTHORIZATION POLICY

POLICY: Anticoagulants – Savaysa Prior Authorization Policy

- Savaysa® (edoxaban tablets – Daiichi Sankyo)

REVIEW DATE: 01/11/2023

OVERVIEW

Savaysa, a Factor Xa inhibitor, is indicated for the following uses:¹

- **Non-valvular atrial fibrillation**, to reduce the risk of stroke and systemic embolism.
- **Treatment of deep vein thrombosis (DVT) and pulmonary embolism (PE)**, following 5 to 10 days of initial therapy with a parenteral anticoagulant.

Savaysa has a unique Boxed Warning regarding reduced efficacy in non-valvular atrial fibrillation in patients with a creatinine clearance > 95 mL/min; Savaysa should be avoided in these individuals.¹ Safety and effectiveness of Savaysa in pediatric patients have not been established.

Guidelines

Guidelines are available which support the use of direct oral anticoagulants (DOACs) in their commonly used clinical settings, such as DVT/PE²⁻⁵ and atrial fibrillation^{6,7}. In patients who are eligible for a DOAC, these are generally preferred over vitamin K antagonists (e.g., warfarin). It is noted that in the randomized trials in atrial fibrillation, DOACs were consistently at least non-inferior to warfarin regarding the composite of stroke or systemic embolism and were associated with lower risk of serious bleeding.⁷

Anticoagulants and Coronavirus Disease 19 (COVID-19)

Several clinical practice guidelines have been published with regard to use of anticoagulant therapy in the management of COVID-19. Per National Institutes of Health treatment guidelines regarding antithrombotic therapy in patients with COVID-19 (updated December 28, 2022), hospitalized patients with COVID-19 should not be routinely discharged from the hospital while on venous thromboembolism (VTE) prophylaxis.⁸ For patients at low risk for bleeding and high risk for VTE, continuing anticoagulation with an FDA-approved regimen for extended VTE prophylaxis may be considered, as per protocols for patients without COVID-19. Of note, Xarelto® (rivaroxaban tablets and oral suspension) is FDA-approved for prophylaxis of VTE in acutely ill medical patients; Savaysa is not indicated in this setting. Other guidelines have similar recommendations.⁹⁻¹¹

Other Uses with Supportive Evidence

Savaysa has data for prophylaxis of VTE after hip replacement surgery.¹² Although data are not robust regarding use of DOACs in other off-label thromboembolic-related conditions, CHEST guidelines (2012) suggest anticoagulation for certain patients with superficial vein thrombosis, symptomatic splanchnic thromboses (portal, mesenteric, and/or splenic vein), or symptomatic hepatic vein thrombosis.² The guidelines acknowledge the limited available data in these settings, and all are given Grade 2C recommendations (weak recommendation, low-quality evidence). The 2016 CHEST guideline update did not address these conditions or comment on the role of DOACs.³ The choice of anticoagulant is often individualized based on patient-specific factors; therefore, for certain patients, DOAC use may be considered in practice. Evidence for DOACs is limited for off-label scenarios; in general, agents

such as vitamin K antagonists (e.g., warfarin) and low molecular weight heparin have more clinical experience in these settings.

POLICY STATEMENT

Prior Authorization is recommended for prescription benefit coverage of Savaysa. All approvals are provided for the duration noted below. In cases where the approval is authorized in months, 1 month is equal to 30 days.

Automation: None.

RECOMMENDED AUTHORIZATION CRITERIA

Coverage of Savaysa is recommended in those who meet one of the following criteria:

FDA-Approved Indications

- 1. Atrial Fibrillation (or Atrial Flutter).** Approve for 1 year if the patient meets both of the following criteria (A and B):
 - A) Patient is ≥ 18 years of age; AND
 - B) Patient has an estimated creatinine clearance ≤ 95 mL/min.
- 2. Deep Vein Thrombosis or Pulmonary Embolism, Treatment.** Approve for 1 year if the patient is ≥ 18 years of age.

Other Uses with Supportive Evidence

- 3. Deep Vein Thrombosis in a Patient Undergoing Hip Replacement Surgery, Prophylaxis.** Approve for 60 days if the patient is ≥ 18 years of age.
- 4. Treatment or Prevention of Other Thromboembolic-Related Conditions.** Approve for 6 months if the patient meets both of the following criteria (A and B):

Note: Examples of other thromboembolic-related conditions include superficial vein thrombosis, splanchnic vein thrombosis, hepatic vein thrombosis, or prophylaxis of venous thromboembolism in a high-risk patient.

 - A) Patient is ≥ 18 years of age; AND
 - B) Patient meets one of the following criteria (i or ii):
 - i. Patient has tried warfarin, fondaparinux, or a low molecular weight heparin product (e.g., enoxaparin, Fragmin [dalteparin injection]); OR
Note: A patient who has tried Eliquis (apixaban tablets), Xarelto (rivaroxaban tablets), or Pradaxa (dabigatran capsules) is not required to try warfarin, fondaparinux, or a low molecular weight heparin.
 - ii. Patient has been started on Savaysa for the treatment of an acute thromboembolic condition.

CONDITIONS NOT RECOMMENDED FOR APPROVAL

Coverage of Savaysa is not recommended in the following situations:

- 1. Venous Thromboembolism in an Acutely Ill Medical Patient, Prophylaxis.** (Note: This includes post-discharge thromboprophylaxis for a patient hospitalized with coronavirus disease 19 [COVID-19]). Xarelto is labeled for prophylaxis of venous thromboembolism in acutely ill medical patients and is supported in clinical practice guidelines, including guidelines which address prophylaxis of venous thromboembolism in COVID-19 patients.⁷⁻⁹

2. Coverage is not recommended for circumstances not listed in the Recommended Authorization Criteria. Criteria will be updated as new published data are available.

REFERENCES

1. Savaysa® tablets [prescribing information]. Basking Ridge, NJ: Daiichi Sankyo; April 2021.
2. Guyatt GH, Akl EA, Crowther M, et al, for the American College of Chest Physicians Antithrombotic Therapy and Prevention of Thrombosis Panel. Executive summary: antithrombotic therapy and prevention of thrombosis, 9th ed: American College of Chest Physicians Evidence-Based Clinical Practice Guidelines. *Chest*. 2012;141:7S-47S.
3. Kearon C, Akl EA, Ornelas J, et al. Antithrombotic therapy for VTE disease: CHEST Guideline and Expert Panel Report. *Chest*. 2016;149(2):315-352.
4. The NCCN Cancer-Associated Venous Thromboembolic Disease Clinical Practice Guidelines in Oncology (version 1.2022 – March 11, 2022). © 2022 National Comprehensive Cancer Network. Available at: <http://www.nccn.org>. Accessed on January 5, 2023.
5. Ortel TL, Neumann I, Ageno W, Beyth R, et al. American Society of Hematology 2020 guidelines for management of venous thromboembolism: treatment of deep vein thrombosis and pulmonary embolism. *Blood Adv*. 2020 Oct 13;4(19):4693-4738.
6. Lip G, Banerjee A, Boriani G, et al. Antithrombotic therapy for atrial fibrillation: CHEST guideline and expert panel report. *Chest*. 2018;154(5):1121-1201.
7. January CT, Wann LS, Calkins H, et al. 2019 AHA/ACC/HRS focused update of the 2014 AHA/ACC/HRS guideline for the management of patients with atrial fibrillation: a report of the American College of Cardiology/American Heart Association Task Force on Clinical Practice Guidelines and the Heart Rhythm Society. *J Am Coll Cardiol*. 2019;74(1):104-132.
8. COVID-19 Treatment Guidelines Panel. Coronavirus Disease 2019 (COVID-19) Treatment Guidelines. National Institutes of Health. Updated December 28, 2022. Available at: <https://www.covid19treatmentguidelines.nih.gov/>. Accessed on January 5, 2023.
9. Moores LK, Tritschler T, Brosnahan S, et al. Prevention, diagnosis, and treatment of VTE in patients with Coronavirus Disease 2019: CHEST Guideline and Expert Panel Report. *Chest*. 2020 Sep;158(3):1143-1163.
10. Spyropoulos AC, Levy JH, Ageno W, et al. Scientific and Standardization Committee communication: Clinical guidance on the diagnosis, prevention, and treatment of venous thromboembolism in hospitalized patients with COVID-19. *J Thromb Haemost*. 2020; 18: 1859– 1865.
11. Barnes GD, Burnett A, Allen A, et al. Thromboembolism and anticoagulant therapy during the COVID-19 pandemic: interim clinical guidance from the anticoagulation forum. *J Thromb Thrombolysis*. 2020 Jul;50(1):72-81.
12. Raskob G, Cohen AT, Eriksson BI, et al. Oral direct factor Xa inhibition with edoxaban for thromboprophylaxis after elective total hip replacement. A randomized double-blind, dose-response study. *Thromb Haemost*. 2010;104(3):642-649.