

PRIOR AUTHORIZATION POLICY

POLICY: Enzyme Replacement Therapy – Fabrazyme Prior Authorization Policy

- Fabrazyme® (agalsidase intravenous infusion – Genzyme)

REVIEW DATE: 04/06/2022

OVERVIEW

Fabrazyme, a human α -galactosidase A (α -Gal), is indicated for use in patients with **Fabry disease**.¹ It is the same amino acid sequence as the native enzyme and is produced in Chinese hamster ovary cells via recombinant DNA technology. Fabrazyme catalyzes the breakdown of globotriaosylceramide (GL-3) and other α -galactyl-terminated neutral glycosphingolipids to ceramide and galactose and reduces the deposition of GL-3 in the capillary endothelium of the kidney and certain other cell types.

Disease Overview

Fabry disease is a rare inherited X-linked lysosomal storage disorder due to absent or significantly reduced α -Gal activity leading to the accumulation of GL-3 in a wide variety of cells throughout the body.²⁻⁴ The accumulation of GL-3 leads to progressive multisystem disease, primarily impacting the kidney, heart and nervous system.^{3,4} The incidence of Fabry disease is estimated to be about 1:117,000 live male births.² Fabry disease can be divided into two phenotypes. A severe, classical phenotype typically occurs in men without α -Gal activity, whereas a generally milder non-classical phenotype is found in men and women with some residual α -Gal activity.^{2,3} The diagnosis of Fabry disease can be confirmed in males by demonstrating a deficiency in α -Gal activity, and in all patients by identifying a Fabry disease causing gene mutation.⁴ Long-term consequences of Fabry disease include hypertrophic cardiomyopathy, arrhythmias, renal failure, and stroke.³ The kidney disease that occurs in Fabry disease is associated with progressive proteinuria and a decline in glomerular filtration rate, which over time, leads to end-stage renal disease requiring dialysis and ultimately, kidney transplantation.² Treatment with Fabrazyme reduces the accumulation of GL-3 in the kidney (and in other organs), with the goal of stopping or slowing the decline in kidney function.

POLICY STATEMENT

Prior Authorization is recommended for prescription benefit coverage of Fabrazyme. All approvals are provided for the duration noted below. Because of the specialized skills required for evaluation and diagnosis of patients treated with Fabrazyme as well as the monitoring required for adverse events and long-term efficacy, approval requires Fabrazyme to be prescribed by or in consultation with a physician who specializes in the condition being treated.

Automation: None.

RECOMMENDED AUTHORIZATION CRITERIA

Coverage of Fabrazyme is recommended in those who meet the following criteria:

FDA-Approved Indication

1. **Fabry Disease.** Approve for 1 year if the patient meets the following criteria (A and B):
 - A) The diagnosis is established by one of the following (i or ii):
 - i. Patient has a laboratory test demonstrating deficient α -galactosidase A activity in leukocytes or fibroblasts; OR
 - ii. Patient has a molecular genetic test demonstrating mutations in the galactosidase alpha gene;
AND
 - B) Fabrazyme is prescribed by or in consultation with a geneticist, endocrinologist, a metabolic disorder sub-specialist, or a physician who specializes in the treatment of lysosomal storage disorders.

CONDITIONS NOT RECOMMENDED FOR APPROVAL

Coverage of Fabrazyme is not recommended in the following situations:

1. Coverage is not recommended for circumstances not listed in the Recommended Authorization Criteria. Criteria will be updated as new published data are available.

REFERENCES

1. Fabrazyme® intravenous infusion [prescribing information]. Cambridge, MA: Genzyme; August 2021.
2. Schiffmann R. Fabry Disease. *Handb Clin Neurol.* 2015;132:231-248.
3. Arends M, Wanner C, Hughes D, et al. Characterization of Classical and Nonclassical Fabry Disease: A Multinational Study. *J Am Soc Nephrol.* 2017;28:1631-1641.
4. Laney DA, Bennett RL, Clarke V, et al. Fabry Disease Practice Guidelines: Recommendations of the National Society of Genetic Counselors. *J Genet Counsel.* 2013;22:555-564.