

PRIOR AUTHORIZATION POLICY

POLICY: Inflammatory Conditions – Sotyktu Prior Authorization Policy

- Sotyktu™ (deucravacitinib tablets – Bristol Myers Squibb)

REVIEW DATE: 09/14/2022

OVERVIEW

Sotyktu, a tyrosine kinase 2 (TYK2) inhibitor, is indicated for treatment of moderate to severe **plaque psoriasis** in adults who are candidates for systemic therapy or phototherapy.¹ Limitation of use: Sotyktu is not recommended in combination with potent immunosuppressants.

Guidelines

Guidelines have not been updated to address Sotyktu. Joint guidelines from the American Academy of Dermatology and National Psoriasis Medical Board (2019) have been published for management of psoriasis with biologics.² These guidelines list all the biologics approved at the time of publication as agents that may be used as monotherapy for adults with moderate to severe psoriasis. Guidelines from the European Dermatology Forum (2015) recommend biologics (i.e., etanercept, adalimumab, infliximab, Stelara [ustekinumab subcutaneous injection]) as second-line therapy for induction and long-term treatment if phototherapy and conventional systemic agents have failed, are contraindicated, or are not tolerated.³

POLICY STATEMENT

Prior Authorization is recommended for prescription benefit coverage of Sotyktu. Because of the specialized skills required for evaluation and diagnosis of patients treated with Sotyktu as well as the monitoring required for adverse events and long-term efficacy, initial approval requires Sotyktu to be prescribed by or in consultation with a physician who specializes in the condition being treated. All approvals are for the duration noted below. In cases where the approval is authorized in months, 1 month is equal to 30 days.

Automation: None.

RECOMMENDED AUTHORIZATION CRITERIA

Coverage of Sotyktu is recommended in those who meet the following criteria:

FDA-Approved Indication

- 1. Plaque Psoriasis.** Approve for the duration noted if the patient meets ONE of the following (A or B):
 - A) Initial Therapy.** Approve for 3 months if the patient meets ALL of the following criteria (i, ii, and iii):
 - i.** Patient is ≥ 18 years of age; AND
 - ii.** Patient meets ONE of the following (a or b):
 - a)** Patient has tried at least one traditional systemic agent for psoriasis for at least 3 months, unless intolerant; OR

Note: Examples of one traditional systemic agent include methotrexate, cyclosporine, acitretin tablets, or psoralen plus ultraviolet A light (PUVA). An exception to the requirement for a trial of one traditional systemic agent for psoriasis can be made if the

09/14/2022

© 2022. All Rights Reserved.

This document is confidential and proprietary. Unauthorized use and distribution are prohibited.

patient has already had a 3-month trial or previous intolerance to at least one biologic other than the requested drug. A biosimilar of the requested biologic does not count. Refer to [Appendix](#) for examples of biologics used for plaque psoriasis. A patient who has already tried a biologic for psoriasis is not required to “step back” and try a traditional systemic agent for psoriasis.

- b) Patient has a contraindication to methotrexate, as determined by the prescriber; AND
 - iii. The medication is prescribed by or in consultation with a dermatologist.
- B) Patient is Currently Receiving Sotyktu.** Approve for 1 year meets ALL of the following (i, ii, and iii):
- i. Patient has been established on therapy for at least 90 days; AND
Note: A patient who has received < 90 days of therapy or who is restarting therapy is reviewed under criterion A (Initial Therapy).
 - ii. Patient experienced a beneficial clinical response, defined as improvement from baseline (prior to initiating the requested drug) in at least one of the following: estimated body surface area, erythema, induration/thickness, and/or scale of areas affected by psoriasis; AND
 - iii. Compared with baseline (prior to initiating the requested drug), patient experienced an improvement in at least one symptom, such as decreased pain, itching, and/or burning.

CONDITIONS NOT RECOMMENDED FOR APPROVAL

Coverage of Sotyktu is not recommended in the following situations:

- 1. Concurrent Use with other Biologics or with Targeted Synthetic Disease-Modifying Antirheumatic Drugs (DMARDs).** Data are lacking evaluating concomitant use of Sotyktu with another biologic or with a targeted synthetic DMARD for an inflammatory condition (see [Appendix](#) for examples). Combination therapy with biologics and/or biologics + targeted synthetic DMARDs has a potential for a higher rate of adverse effects and lack controlled trial data in support of additive efficacy.⁴
Note: This does NOT exclude the use of methotrexate (a traditional systemic agent used to treat psoriasis) in combination with Sotyktu.
- 2. Concurrent use with Other Potent Immunosuppressants, Including Methotrexate.**¹ Co-administration with other potent immunosuppressive drugs has the risk of added immunosuppression and has not been evaluated.
- 3.** Coverage is not recommended for circumstances not listed in the Recommended Authorization Criteria. Criteria will be updated as new published data are available.

REFERENCES

1. Sotyktu™ tablets [prescribing information]. Princeton, NJ: Bristol Myers Squibb; September 2022.
2. Menter A, Strober BE, Kaplan DH, et al. Joint AAD-NPF guidelines of care for the management and treatment of psoriasis with biologics. *J Am Acad Dermatol.* 2019;80(4):1029-1072.
3. Nast A, Gisondi P, Ormerod AD, et al. European S3-Guidelines on the systemic treatment of psoriasis vulgaris – Update 2015 – Short version – EDF in cooperation with EADV and IPC. *J Eur Acad Dermatol Venereol.* 2015;29(12):2277-2294.

Type of Revision	Summary of Changes	Review Date
New Policy	--	09/14/2022

APPENDIX

* Not an all-inclusive list of indications (e.g., oncology indications and rare inflammatory conditions are not listed). Refer to the prescribing information for the respective agent for FDA-approved indications; SC – Subcutaneous; TNF – Tumor necrosis factor; AS – Ankylosing spondylitis; CD – Crohn’s disease; JIA – Juvenile idiopathic arthritis; PsO – Plaque psoriasis; PsA – Psoriatic arthritis; RA – Rheumatoid arthritis; UC – Ulcerative colitis; nr-axSpA – Non-radiographic axial spondyloarthritis; IV – Intravenous, PJIA – Polyarticular juvenile idiopathic arthritis; IL – Interleukin; SJIA – Systemic juvenile idiopathic arthritis; ^ Off-label use of Kineret in JIA supported in guidelines; ERA – Enthesitis-related arthritis; DMARD – Disease-modifying antirheumatic drug; PDE4 – Phosphodiesterase 4; JAK – Janus kinase; AD – Atopic dermatitis; TYK2 – Tyrosine kinase 2.