

PRIOR AUTHORIZATION POLICY

POLICY: Oncology (Injectable – CAR-T) – Tecartus Prior Authorization Policy

- Tecartus® (brexucabtagene autoleucl intravenous infusion – Kite Pharma)

REVIEW DATE: 08/21/2024

OVERVIEW

Tecartus, a CD19-directed genetically modified autologous T cell immunotherapy, is indicated for the treatment of adults with relapsed or refractory:¹

- **B-cell precursor acute lymphoblastic leukemia.**
- **Mantle cell lymphoma.**

Tecartus is supplied in infusion bag(s) containing frozen suspension of genetically modified autologous T cells in human serum albumin.¹ Each bag is supplied in a metal cassette stored in the vapor phase of liquid nitrogen. Store Tecartus frozen in the vapor phase of liquid nitrogen and thaw prior to administration.

Guidelines

Tecartus is addressed in National Comprehensive Cancer Network guidelines:

- **Acute lymphoblastic leukemia:** Guidelines (version 2.2024 – July 19, 2024) recommend Tecartus for the treatment of relapsed or refractory B-cell precursor acute lymphoblastic leukemia.^{3,4}
- **B-cell lymphomas:** Guidelines (version 2.2024 – April 30, 2024) recommend Tecartus for the second-line and subsequent treatment of relapsed or refractory mantle cell lymphoma, following treatment with Bruton tyrosine kinase inhibitor therapy.^{2,3}

Safety

Tecartus has a Boxed Warning regarding cytokine release syndrome, neurological toxicities, and T-cell malignancies.¹ Due to these risks, Tecartus is only available through a restricted program under a Risk Evaluation and Mitigation Strategy (REMS) called the Tecartus REMS.

POLICY STATEMENT

Prior Authorization is recommended for prescription benefit coverage of Tecartus. All approvals for therapy are provided for the approval duration noted below. Because of the specialized skills required for evaluation and diagnosis of patients treated with Tecartus as well as the monitoring required for adverse events and long-term efficacy, approval requires Tecartus to be prescribed by or in consultation with a physician who specializes in the condition being treated. The approval duration is 6 months to allow for an adequate time frame to prepare and administer 1 dose of therapy.

Automation: None.

RECOMMENDED AUTHORIZATION CRITERIA

Coverage of Tecartus is recommended in those who meet one of the following criteria:

FDA-Approved Indications

08/21/2024

© 2024. All Rights Reserved.

This document is confidential and proprietary. Unauthorized use and distribution are prohibited.

1. **Acute Lymphoblastic Leukemia.** Approve a single dose if the patient meets ALL of the following (A, B, C, D, E, and F):
 - A) Patient is \geq 18 years of age; AND
 - B) Patient has B-cell precursor disease; AND
 - C) Patient has relapsed or refractory disease; AND
 - D) Patient received or plans to receive lymphodepleting chemotherapy prior to Tecartus infusion; AND
 - E) Patient has not been previously treated with CAR-T therapy; AND
Note: Examples of CAR-T therapy include Tecartus, Breyanzi (lisocabtagene maraleucel intravenous infusion), Kymriah (tisagenlecleucel intravenous infusion), Yescarta (axicabtagene intravenous infusion) and Abecma (idecabtagene vicleucel intravenous infusion).
 - F) Tecartus is prescribed by or in consultation with an oncologist.

2. **Mantle Cell Lymphoma.** Approve a single dose if the patient meets ALL of the following (A, B, C, D, and E):
 - A) Patient is \geq 18 years of age; AND
 - B) Patient has relapsed or refractory disease; AND
 - C) Patient received or plans to receive lymphodepleting chemotherapy prior to Tecartus infusion; AND
 - D) Patient has not been previously treated with CAR-T therapy; AND
Note: Examples of CAR-T therapy include Tecartus, Breyanzi (lisocabtagene maraleucel intravenous infusion), Kymriah (tisagenlecleucel intravenous infusion), Yescarta (axicabtagene intravenous infusion) and Abecma (idecabtagene vicleucel intravenous infusion).
 - E) Tecartus is prescribed by or in consultation with an oncologist.

CONDITIONS NOT RECOMMENDED FOR APPROVAL

Coverage of Tecartus is not recommended in the following situations:

1. Coverage is not recommended for circumstances not listed in the Recommended Authorization Criteria. Criteria will be updated as new published data are available.

REFERENCES

1. Tecartus® intravenous infusion [prescribing information]. Santa Monica, CA: Kite Pharma; June 2024.
2. The NCCN B-Cell Lymphomas Clinical Practice Guidelines in Oncology (version 2.2024 – April 30, 2024). © 2024 National Comprehensive Cancer Network. Available at: <http://www.nccn.org>. Accessed on August 13, 2024.
3. The NCCN Drugs and Biologics Compendium. © 2024 National Comprehensive Cancer Network. Available at: <http://www.nccn.org>. Accessed on August 13, 2024. Search term: brexucabtagene.
4. The NCCN Acute Lymphoblastic Leukemia Clinical Practice Guidelines in Oncology (version 2.2024 – July 19, 2024). © 2024 National Comprehensive Cancer Network. Available at: <http://www.nccn.org>. Accessed on August 13, 2024.