

## PRIOR AUTHORIZATION POLICY

**POLICY:** Oncology (Injectable) – Besremi Prior Authorization Policy

- Besremi® (ropeginterferon alfa-2b-njft subcutaneous injection – PharmaEssentia)

**REVIEW DATE:** 11/16/2022

---

### OVERVIEW

Besremi, an interferon alfa, is indicated for treatment of adults with **polycythemia vera**.<sup>1</sup>

### Guidelines

The National Comprehensive Cancer Network guidelines for myeloproliferative neoplasms (version 3.2022 – August 11, 2022) discuss therapies for polycythemia vera.<sup>2</sup> In low-risk patients, management of cardiovascular risk factors, low-dose aspirin (81 to 100 mg/day), and phlebotomy to maintain hematocrit < 45% are recommended (category 2A for all). Besremi is listed as another recommended regimen (category 2B). A footnote comments that cytoreductive therapy is not recommended for initial treatment. In high-risk patients, preferred regimens for cytoreductive therapy include hydroxyurea or Pegasys® (peginterferon alfa-2a subcutaneous injection) [category 2A for both]. Besremi is another recommended regimen in this setting (category 2A). A footnote comments that Pegasys can be considered for younger patients, pregnant patients, or patients who defer hydroxyurea or Besremi.

### POLICY STATEMENT

Prior Authorization is recommended for prescription benefit coverage of Besremi. All approvals are provided for the duration noted below. Because of the specialized skills required for evaluation and diagnosis of patients treated with Besremi as well as the monitoring required for adverse events and long-term efficacy, approval requires Besremi to be prescribed by or in consultation with a physician who specializes in the condition being treated.

**Automation:** None.

### RECOMMENDED AUTHORIZATION CRITERIA

Coverage of Besremi is recommended in those who meet the following criteria:

#### FDA-Approved Indication

1. **Polycythemia Vera.** Approve for 1 year the patient meets the following criteria (A and B):
  - A) Patient is  $\geq 18$  years of age; AND
  - B) Besremi is prescribed by or in consultation with an oncologist.

### CONDITIONS NOT RECOMMENDED FOR APPROVAL

Coverage of Besremi is not recommended in the following situations:

1. **Concomitant Use with Other Interferon Products.** Besremi was not studied in combination with other interferon products; concomitant use would be expected to result in increased toxicity.  
Note: An example of an interferon product is Pegasys® (peginterferon alfa-2a subcutaneous injection).

11/16/2022

© 2022. All Rights Reserved.

This document is confidential and proprietary. Unauthorized use and distribution are prohibited.

2. **Hepatitis B Virus.** Besremi is not indicated for hepatitis B.<sup>1</sup> Pegylated interferons are recommended in American Association for the Study of Liver Diseases (AASLD) guidelines for chronic hepatitis B (updated 2018).<sup>3</sup> Phase I/II data suggest similar efficacy between Besremi and Pegasys for chronic hepatitis B; however, further data are needed.<sup>4</sup>
3. **Hepatitis C Virus.** Besremi is not indicated for hepatitis C.<sup>1</sup> Pegasys, another pegylated interferon, is indicated for the treatment of chronic hepatitis C. However, peginterferons are no longer addressed by the AASLD recommendations for testing, managing, and treating HCV (updated October 24, 2022).<sup>5</sup>
4. Coverage is not recommended for circumstances not listed in the Recommended Authorization Criteria. Criteria will be updated as new published data are available.

## REFERENCES

1. Besremi<sup>®</sup> subcutaneous injection [prescribing information]. Burlington, MA: PharmaEssentia; November 2021.
2. The NCCN Myeloproliferative Neoplasms Clinical Practice Guidelines in Oncology (version 3.2022 – August 11, 2022). © 2022 National Comprehensive Cancer Network. Available at: <http://www.nccn.org>. Accessed on November 14, 2022.
3. Terrault NA, Lok ASF, McMahon BJ, et al. Update on prevention, diagnosis, and treatment of chronic hepatitis B: AASLD 2018 hepatitis B guidance. *Hepatology*. 2018 Apr;67(4):1560-1599.
4. Huang YW, Hsu CW, Lu SN, et al. Ropeginterferon alfa-2b every 2 weeks as a novel pegylated interferon for patients with chronic hepatitis B. *Hepatol Int*. 2020 Dec;14(6):997-1008.
5. American Association for the Study of Liver Diseases and the Infectious Diseases Society of America. Testing, managing, and treating hepatitis C. Updated October 24, 2022. Available at: <http://www.hcvguidelines.org>. Accessed on: November 14, 2022.

NA – Not applicable.