

## PRIOR AUTHORIZATION POLICY

**POLICY:** Oncology (Injectable) – Cyramza Prior Authorization Policy

- Cyramza® (ramucirumab intravenous infusion – Eli Lilly)

**REVIEW DATE:** 06/15/2022

---

### OVERVIEW

Cyramza, a human vascular endothelial growth factor receptor 2 (VEGFR2) antagonist, is indicated for the following:<sup>1</sup>

- **Colorectal cancer**, metastatic, in combination with FOLFIRI (irinotecan, leucovorin, and 5-fluorouracil [5-FU]) for the treatment of patients with disease progression on or after prior therapy with bevacizumab, oxaliplatin, and a fluoropyrimidine.
- **Gastric or gastroesophageal junction adenocarcinoma**, as a single agent or in combination with paclitaxel for the treatment of patients with advanced or metastatic disease with disease progression on or after prior fluoropyrimidine- or platinum-containing chemotherapy.
- **Hepatocellular carcinoma**, as a single agent in patients who have an alpha fetoprotein of  $\geq 400$  ng/mL and have been treated with Nexavar® (sorafenib tablets).
- **Non-small cell lung cancer (NSCLC)**, metastatic, in combination with docetaxel for the treatment of patients with disease progression on or after platinum-based chemotherapy. Patients with epidermal growth factor receptor (*EGFR*) or anaplastic lymphoma kinase (*ALK*) genomic tumor aberrations should have disease progression on FDA-approved therapy for these aberrations prior to receiving Cyramza.
- **NSCLC**, metastatic, in combination with erlotinib for the first-line treatment of NSCLC with *EGFR* exon 19 deletions or exon 21 (L858R) mutations.

### Guidelines

Cyramza is addressed in National Comprehensive Cancer Network (NCCN) guidelines:

- **Colon cancer** (version 1.2022 – February 25, 2022) and **rectal cancer** (version 1.2022 – February 25, 2022): Guidelines recommend Cyramza as primary therapy and subsequent therapy for patients with unresectable advanced or metastatic disease, and as adjuvant treatment for unresectable metachronous metastases that converted to resectable disease after primary treatment, in combination with either irinotecan or FOLFIRI.<sup>2-4</sup>
- **Gastric cancer** (version 2.2022 – January 11, 2022) and **esophageal and esophagogastric junction cancers** (version 2.2022 – February 11, 2022): Guidelines recommend Cyramza as palliative treatment for patients who are not surgical candidates or have unresectable locally advanced, recurrent, or metastatic disease.<sup>4-6</sup>
- **Hepatobiliary cancers** (version 1.2022 – March 29, 2022): Guidelines recommend Cyramza as a single agent for the treatment of patients with progressive disease with an alpha fetoprotein  $\geq 400$  ng/mL and Child-Pugh Class A only.<sup>4,8</sup>
- **NSCLC** (version 3.2022 – March 16, 2022): Guidelines recommend Cyramza as subsequent therapy in combination with docetaxel for metastatic disease for patients who have not previously received docetaxel either following progression on initial cytotoxic therapy or for further progression on a systemic immune checkpoint inhibitor or other systemic therapy.<sup>4,7</sup> Cyramza is also recommended in combination with erlotinib for patients with *EGFR* exon 19 deletion or L858R mutation positive, recurrent, advanced, or metastatic disease as first-line therapy or as continuation therapy following disease progression on Cyramza and erlotinib.

### POLICY STATEMENT

06/15/2022

© 2022. All Rights Reserved.

This document is confidential and proprietary. Unauthorized use and distribution are prohibited.

Prior Authorization is recommended for prescription benefit coverage of Cyramza. All approvals are provided for the duration noted below. Because of the specialized skills required for evaluation and diagnosis of patients treated with Cyramza as well as the monitoring required for adverse events and long-term efficacy, approval requires Cyramza to be prescribed by or in consultation with a physician who specializes in the condition being treated.

**Automation:** None.

## RECOMMENDED AUTHORIZATION CRITERIA

Coverage of Cyramza is recommended in those who meet one of the following criteria:

### FDA-Approved Indications

1. **Colon or Rectal Cancer.** Approve for 1 year if the patient meets ALL of the following criteria (A, B, and C):
  - A) Patient has received both of the following (i and ii):
    - i. Oxaliplatin; AND
    - ii. Fluoropyrimidine (e.g., 5-fluorouracil [5-FU], capecitabine); AND
  - B) Cyramza will be used in combination with one of the following (i or ii):
    - i. Irinotecan; OR
    - ii. FOLFIRI (irinotecan, folinic acid [leucovorin], and 5-fluorouracil [5-FU]); AND
  - C) Cyramza is prescribed by or in consultation with an oncologist.
2. **Gastric, Esophagogastric Junction, or Esophageal Cancer.** Approve for 1 year if the patient meets ALL of the following criteria (A, B, and C):
  - A) Patient meets one of the following criteria (i, ii, or iii):
    - i. Cyramza will be used alone; OR
    - ii. Cyramza will be used in combination with paclitaxel; OR
    - iii. Cyramza will be used in combination with irinotecan; AND
  - B) Patient has received chemotherapy with at least ONE of the following (i or ii):
    - i. 5-Fluorouracil (5-FU) or capecitabine; OR
    - ii. Cisplatin, carboplatin, or oxaliplatin.
  - C) Cyramza is prescribed by or in consultation with an oncologist.
3. **Hepatocellular Carcinoma.** Approve for 1 year if the patient meets ALL of the following criteria (A, B, C, D, and E):
  - A) Patient has been treated with Nexavar (sorafenib tablet); AND
  - B) Cyramza will be used as a single agent; AND
  - C) Patient has an alpha fetoprotein of  $\geq 400$  ng/mL; AND
  - D) Patient has Child-Pugh Class A disease; AND
  - E) Cyramza is prescribed by or in consultation with an oncologist.
4. **Non-Small Cell Lung Cancer.** Approve for 1 year if the patient meets BOTH of the following criteria (A and B):
  - A) Patient meets of the following criteria (i or ii):
    - i. Cyramza will be used as first-line therapy; AND
      - a) Patient has epidermal growth factor receptor (EGFR) exon 19 deletion or L858R mutation positive disease; AND
      - b) Cyramza will be used in combination with erlotinib; OR

- ii. Cyramza will be used as subsequent therapy; AND
  - a) Cyramza will be used in combination with docetaxel intravenous infusion; AND
  - b) Patient has received targeted drug therapy if the patient’s tumor is positive for a targetable mutation; AND

Note: Examples of targetable mutations include sensitizing epidermal growth factor receptor mutation, anaplastic lymphoma kinase fusions.

- B) Cyramza is prescribed by or in consultation with an oncologist.

### CONDITIONS NOT RECOMMENDED FOR APPROVAL

Coverage of Cyramza is not recommended in the following situations:

1. Coverage is not recommended for circumstances not listed in the Recommended Authorization Criteria. Criteria will be updated as new published data are available.

### REFERENCES

1. Cyramza® intravenous infusion [prescribing information]. Indianapolis, IN: Eli Lilly; March 2022.
2. The NCCN Colon Cancer Clinical Practice Guidelines in Oncology (version 1.2022 – February 25, 2022). © 2022 National Comprehensive Cancer Network. Available at: <http://www.nccn.org>. Accessed on June 8, 2022.
3. The NCCN Rectal Cancer Clinical Practice Guidelines in Oncology (version 2.2022 – February 25, 2022). © 2022 National Comprehensive Cancer Network. Available at: <http://www.nccn.org>. Accessed on June 8, 2022.
4. The NCCN Drugs & Biologics Compendium. © 2022 National Comprehensive Cancer Network. Available at: <http://www.nccn.org>. Accessed on June 8, 2022. Search term: ramucirumab.
5. The NCCN Gastric Cancer Clinical Practice Guidelines in Oncology (version 2.2022 – January 11, 2022). © 2022 National Comprehensive Cancer Network. Available at: <http://www.nccn.org>. Accessed on June 8, 2022.
6. The NCCN Esophageal and Esophagogastric Junction Cancers Clinical Practice Guidelines in Oncology (version 2.2022 – February 11, 2022). © 2022 National Comprehensive Cancer Network. Available at: <http://www.nccn.org>. Accessed on June 8, 2022.
7. The NCCN Non-Small Cell Lung Cancer Clinical Practice Guidelines in Oncology (version 3.2022 – March 16, 2022). © 2022 National Comprehensive Cancer Network. Available at: <http://www.nccn.org>. Accessed on June 8, 2022.
8. The NCCN Hepatobiliary Cancers Clinical Practice Guidelines in Oncology (version 1.2022 – March 29, 2022). © 2022 National Comprehensive Cancer Network. Available at: <http://www.nccn.org>. Accessed on June 8, 2022.