

PRIOR AUTHORIZATION POLICY

- POLICY:** Oncology (Injectable) – Topotecan Products Prior Authorization Policy
- Hycamtin® (topotecan capsule – Novartis)
 - Topotecan intravenous infusion (Hycamtin® – Novartis, generics)

REVIEW DATE: 01/18/2023

OVERVIEW

Topotecan injection, a topoisomerase inhibitor, is indicated for the treatment of patients with:

- **Cervical cancer**, stage IV-B, recurrent, or persistent disease which is not amenable to curative treatment, in combination with cisplatin.
- **Metastatic ovarian cancer**, after disease progression on or after initial or subsequent chemotherapy, as a single agent.
- **Small cell lung cancer (SCLC)**, platinum-sensitive disease who progressed at least 60 days after initiation of first-line chemotherapy, as a single agent.¹

Guidelines

Topotecan is included in a variety of National Comprehensive Cancer Network (NCCN) guidelines:

- **Bone cancer** (version 2.2023 – September 28, 2022) clinical practice guidelines recommend topotecan in combination with cyclophosphamide, as second-line therapy for patients with relapsed/refractory, or metastatic osteosarcoma and Ewing sarcoma (both category 2A), and dedifferentiated chondrosarcoma, high-grade undifferentiated pleomorphic sarcoma, and mesenchymal chondrosarcoma (category 2B).^{2,7}
- **Central nervous system cancers** (version 2.2022 – September 29, 2022) clinical practice guidelines recommend topotecan as a single agent for the treatment of brain metastases in patients with small cell lung cancer.^{2,8} In addition, the guidelines recommend intra-cerebrospinal fluid topotecan for the treatment of leptomeningeal metastases.
- **Cervical cancer** (version 1.2023 – January 6, 2023) clinical practice guidelines recommend topotecan as first-line, second-line, or subsequent therapy for patients with local/regional recurrence, stage IV-B disease, or distant metastases in combination with paclitaxel and bevacizumab (category 1), or in combination with paclitaxel or cisplatin (category 2A); or as a single agent in second-line and subsequent therapy.^{2,5} It is also recommended as first-line, second-line and subsequent therapy for patients with persistent, recurrent, or metastatic small cell neuroendocrine carcinoma of the cervix (NECC) in combination with paclitaxel and bevacizumab. Topotecan can be used in combination with paclitaxel or cisplatin, or as a single agent (category 2A) for second-line or subsequent therapy of NECC.
- **Merkel cell carcinoma** (version 2.2022 – March 24, 2022) clinical practice guidelines recommend topotecan as a treatment option for patients with distant metastatic disease who have contraindications to checkpoint immunotherapy (Bavencio® [avelumab intravenous infusion], Keytruda® [pembrolizumab intravenous infusion], and Opdivo® [nivolumab intravenous infusion]).^{2,9}
- **Ovarian cancer** (version 1.2023 – December 22, 2022) clinical practice guidelines recommend topotecan, as a single agent or in combination with bevacizumab or sorafenib, for the treatment of recurrent or persistent platinum-resistant epithelial ovarian cancer, fallopian tube cancer, and peritoneal cancer.^{2,3} Treatment of clinical relapse is a category 2A recommendation and immediate treatment of biochemical relapse is category 2B recommendation.

01/18/2023

© 2022. All Rights Reserved.

This document is confidential and proprietary. Unauthorized use and distribution are prohibited.

- **SCLC** (version 3.2023 – December 21, 2022) clinical practice guidelines recommend topotecan as a single agent for patients with a performance status of 0-2 and relapse following complete or partial response, or stable disease with initial treatment; or for primary progressive disease.^{2,4}
- **Soft tissue sarcoma** (version 2.2022 – May 17, 2022) clinical practice guidelines recommend topotecan as a single agent or in combination with cyclophosphamide for the treatment of non-pleomorphic rhabdomyosarcoma.^{2,10}
- **Uterine cancer** (version 1.2023 – December 22, 2022) clinical practice guidelines recommend topotecan as a single agent for the treatment of recurrent, metastatic, or high-risk endometrial carcinoma.^{2,6}

POLICY STATEMENT

Prior Authorization is recommended for prescription benefit coverage of topotecan. All approvals are provided for the duration noted below. Because of the specialized skills required for evaluation and diagnosis of patients treated with topotecan as well as the monitoring required for adverse events and long-term efficacy, approval requires topotecan to be prescribed by or in consultation with a physician who specializes in the condition being treated.

Automation: None.

RECOMMENDED AUTHORIZATION CRITERIA

Coverage of topotecan is recommended in those who meet one of the following criteria:

FDA-Approved Indications

- 1. Cervical Cancer.** Approve for 1 year if the patient meets the following criteria (A, B, and C):
 - A) Patient is ≥ 18 years of age; AND
 - B) Patient meets one of the following (i or ii):
 - i. Patient has persistent or recurrent disease; OR
 - ii. Patient has metastatic disease; AND
 - C) Topotecan is prescribed by or in consultation with an oncologist.
- 2. Ovarian, Fallopian Tube, and Primary Peritoneal Cancer.** Approve for 1 year if the patient meets the following criteria (A, B, C, and D):
 - A) Patient is ≥ 18 years of age; AND
 - B) Patient has persistent or recurrent disease; AND
 - C) The cancer is platinum-resistant; AND
 - D) Topotecan is prescribed by or in consultation with an oncologist.
- 3. Small Cell Lung Cancer.** Approve for 1 year if the patient meets the following criteria (A, B, C, and D):
 - A) Patient is ≥ 18 years of age; AND
 - B) Patient meets one of the following (i or ii):
 - i. Patient has relapsed disease; OR
 - ii. Patient has primary progressive disease; AND
 - C) Topotecan will be used as a single agent; AND
 - D) Topotecan is prescribed by or in consultation with an oncologist.

Other Uses with Supportive Evidence

01/18/2023

© 2022. All Rights Reserved.

This document is confidential and proprietary. Unauthorized use and distribution are prohibited.

4. **Bone Cancer.** Approve for 1 year if the patient meets the following criteria (A, B, C, D, and E):
 - A) Patient has one of the following (i, ii, iii, iv, or v):
 - i. Osteosarcoma; OR
 - ii. Ewing sarcoma; OR
 - iii. Dedifferentiated chondrosarcoma; OR
 - iv. High-grade undifferentiated pleomorphic sarcoma; OR
 - v. Mesenchymal chondrosarcoma; AND
 - B) Patient has relapsed, refractory, or metastatic disease; AND
 - C) Topotecan is used second-line; AND
 - D) Topotecan is used in combination with cyclophosphamide; AND
 - E) Topotecan is prescribed by or in consultation with an oncologist.

5. **Brain Metastases.** Approve for 1 year if the patient meets the following criteria (A, B, C, and D):
 - A) Patient is ≥ 18 years of age; AND
 - B) Patient has small cell lung cancer; AND
 - C) Topotecan will be used as a single agent; AND
 - D) Topotecan is prescribed by or in consultation with an oncologist.

6. **Endometrial Carcinoma.** Approve for 1 year if the patient meets the following criteria (A, B, C, and D):
 - A) Patient is ≥ 18 years of age; AND
 - B) Patient has recurrent, metastatic, or high-risk disease; AND
 - C) Topotecan will be used as a single agent; AND
 - D) Topotecan is prescribed by or in consultation with an oncologist.

7. **Leptomeningeal and Spinal Metastases.** Approve for 1 year if the patient meets the following criteria (A, B, and C):
 - A) Patient is ≥ 18 years of age; AND
 - B) Topotecan will be administered intraventricularly; AND
 - C) Topotecan is prescribed by or in consultation with an oncologist.

8. **Merkel Cell Carcinoma.** Approve for 1 year if the patient meets the following criteria (A, B, C, and D):
 - A) Patient is ≥ 18 years of age; AND
 - B) Patient has distant metastatic disease; AND
 - C) Patient has contraindications to checkpoint immunotherapy; AND
Note: Checkpoint immunotherapy includes Bavencio (avelumab intravenous infusion), Keytruda (pembrolizumab intravenous infusion), and Opdivo (nivolumab intravenous infusion).
 - D) Topotecan is prescribed by or in consultation with an oncologist.

9. **Rhabdomyosarcoma.** Approve for 1 year if the patient meets the following criteria (A, B, and C):
 - A) Patient is ≥ 18 years of age; AND
 - B) Patient has non-pleomorphic rhabdomyosarcoma; AND
 - C) Topotecan is prescribed by or in consultation with an oncologist.

CONDITIONS NOT RECOMMENDED FOR APPROVAL

Coverage of topotecan is not recommended in the following situations:

1. Coverage is not recommended for circumstances not listed in the Recommended Authorization Criteria. Criteria will be updated as new published data are available.

REFERENCES

1. Hycamtin® intravenous infusion [prescribing information]. East Hanover, NJ: Novartis; October 2019.
2. The NCCN Drugs & Biologics Compendium. © 2023 National Comprehensive Cancer Network. Available at: <http://www.nccn.org>. Accessed on January 11, 2023. Search term: topotecan.
3. The NCCN Ovarian Cancer Clinical Practice Guidelines (version 1.2023 – December 22, 2022). © 2022 National Comprehensive Cancer Network. Available at: <http://www.nccn.org>. Accessed January 12, 2023.
4. The NCCN Small Cell Lung Cancer Clinical Practice Guidelines (version 3.2023 – December 21, 2022). © 2022 National Comprehensive Cancer Network. Available at: <http://www.nccn.org>. Accessed January 12, 2023.
5. The NCCN Cervical Cancer Clinical Practice Guidelines (version 1.2023 – January 6, 2023). © 2023 National Comprehensive Cancer Network. Available at: <http://www.nccn.org>. Accessed January 12, 2023.
6. The NCCN Uterine Cancer Clinical Practice Guidelines (version 1.2023 – December 22, 2022). © 2022 National Comprehensive Cancer Network. Available at: <http://www.nccn.org>. Accessed January 12, 2023.
7. The NCCN Bone Cancer Clinical Practice Guidelines (version 2.2023 – September 28, 2022). © 2022 National Comprehensive Cancer Network. Available at: <http://www.nccn.org>. Accessed January 12, 2023.
8. The NCCN Central Nervous System Cancers Clinical Practice Guidelines (version 2.2022 – September 29, 2022). © 2022 National Comprehensive Cancer Network. Available at: <http://www.nccn.org>. Accessed January 12, 2023.
9. The NCCN Merkel Cell Carcinoma Clinical Practice Guidelines (version 2.2022 – March 24, 2022). © 2022 National Comprehensive Cancer Network. Available at: <http://www.nccn.org>. Accessed January 11, 2023.
10. The NCCN Soft Tissue Sarcoma Clinical Practice Guidelines (version 2.2022 – May 17, 2022). © 2022 National Comprehensive Cancer Network. Available at: <http://www.nccn.org>. Accessed January 12, 2023.