

PRIOR AUTHORIZATION POLICY

POLICY: Oncology – Bexarotene (Topical) Prior Authorization Policy

- Targretin® (bexarotene 1% gel – Bausch Health, generic)

REVIEW DATE: 11/02/2022; selected revision 01/25/2023

OVERVIEW

Bexarotene gel is indicated for the topical treatment of cutaneous lesions in patients with **cutaneous T-cell lymphoma** (Stage 1A and 1B) who have refractory or persistent disease after other therapies or who have not tolerated other therapies.¹

Guidelines

National Comprehensive Cancer Network (NCCN) Primary Cutaneous Lymphomas guidelines (version 1.2023 – January 5, 2023) recommend topical bexarotene as an option for the treatment of cutaneous lymphomas (e.g., mycosis fungoides, Sézary syndrome, T-cell lymphoma), as initial therapy and for relapsed/refractory cases. NCCN notes there are case reports demonstrating efficacy of topical bexarotene in treating primary cutaneous B-cell lymphomas in children.

POLICY STATEMENT

Prior Authorization is recommended for prescription benefit coverage of bexarotene gel. All approvals are provided for the duration noted below. Because of the specialized skills required for evaluation and diagnosis of patients treated with bexarotene gel as well as the monitoring required for adverse events and long-term efficacy, approval requires bexarotene gel to be prescribed by or in consultation with a physician who specializes in the condition being treated.

Automation: None.

RECOMMENDED AUTHORIZATION CRITERIA

Coverage of Bexarotene gel is recommended in those who meet the following criteria:

FDA-Approved Indication

1. **Cutaneous T-Cell Lymphoma.** Approve for 1 year if the patient meets the following criteria (A and B):
 - A) Patient has cutaneous manifestations/lesions; AND
 - B) The medication is prescribed by or in consultation with an oncologist or a dermatologist.

CONDITIONS NOT RECOMMENDED FOR APPROVAL

Coverage of bexarotene gel is not recommended in the following situations:

1. Coverage is not recommended for circumstances not listed in the Recommended Authorization Criteria. Criteria will be updated as new published data are available.

REFERENCES

1. Targretin® gel [prescribing information]. Bridgewater, NJ: Bausch Health; February 2020.

11/02/2022

© 2022. All Rights Reserved.

This document is confidential and proprietary. Unauthorized use and distribution are prohibited.

2. The NCCN Primary Cutaneous Lymphomas Clinical Practice Guidelines in Oncology (version 2.2022 – June 8, 2022). © 2022 National Comprehensive Cancer Network. Available at: <http://www.nccn.org>. Accessed on October 28, 2022.
3. The NCCN Drugs & Biologics Compendium. © 2023 National Comprehensive Cancer Network. Available at: <http://www.nccn.org>. Accessed on January 23, 2023. Search terms: bexarotene gel.