

PRIOR AUTHORIZATION POLICY

POLICY: Somatostatin Analogs – Lanreotide Products Prior Authorization Policy

- Lanreotide subcutaneous injection – Cipla
- Somatuline[®] Depot (lanreotide subcutaneous injection – Ipsen)

REVIEW DATE: 05/15/2024

OVERVIEW

The lanreotide products are somatostatin analogs indicated for the following uses:^{1,2}

- **Acromegaly**, in patients who have had an inadequate response to surgery and/or radiotherapy, or for whom surgery and/or radiotherapy, is not an option. The goal of treatment in acromegaly is to reduce growth hormone and insulin-like growth factor-1 levels to normal.
- **Gastroenteropancreatic neuroendocrine tumors (GEP-NETs)**, in adult patients with unresectable, well or moderately differentiated, locally advanced or metastatic GEP-NETs to improve progression-free survival.

Additionally, Somatuline Depot is indicated for **carcinoid syndrome**, in adult patients.¹

Guidelines

The National Comprehensive Cancer Network (NCCN) guidelines for **neuroendocrine and adrenal tumors** (version 1.2023 – August 2, 2023) recommend Somatuline Depot for the management of carcinoid syndrome; tumors of the gastrointestinal tract, lung, thymus (carcinoid tumors), and pancreas (including glucagonomas, gastrinomas, VIPomas, insulinomas); pheochromocytomas; and paragangliomas.³ Patients who have local unresectable disease and/or distant metastases and clinically significant tumor burden or progression should be started on therapy with a somatostatin analog to potentially control tumor growth.

POLICY STATEMENT

Prior Authorization is recommended for prescription benefit coverage of lanreotide products. All approvals are provided for the duration noted below. Because of the specialized skills required for evaluation and diagnosis of patients treated with lanreotide products as well as the monitoring required for adverse events and long-term efficacy, approval requires lanreotide products to be prescribed by or in consultation with a physician who specializes in the condition being treated.

Automation: None.

RECOMMENDED AUTHORIZATION CRITERIA

Coverage of lanreotide products is recommended in those who meet one of the following criteria:

I. Coverage of Somatuline Depot is recommended in those who meet one of the following criteria:

FDA-Approved Indications

1. **Acromegaly.** Approve for 1 year if the patient meets ALL of the following (A, B, and C):
 - A) Patient meets ONE of the following (i, ii, or iii):
 - i. Patient has had an inadequate response to surgery and/or radiotherapy; OR

05/15/2024

© 2024. All Rights Reserved.

This document is confidential and proprietary. Unauthorized use and distribution are prohibited.

- ii. Patient is NOT an appropriate candidate for surgery and/or radiotherapy; OR
 - iii. Patient is experiencing negative effects due to tumor size (e.g., optic nerve compression); AND
 - B) Patient has (or had) a pre-treatment (baseline) insulin-like growth factor-1 (IGF-1) level above the upper limit of normal (ULN) based on age and gender for the reporting laboratory; AND
Note: Pre-treatment (baseline) refers to the IGF-1 level prior to the initiation of any somatostatin analog (e.g., Mycapssa [octreotide delayed-release capsules], an octreotide acetate injection product [e.g., Bynfezia Pen, Sandostatin {generics}, Sandostatin LAR Depot], Signifor LAR [pasireotide injection], Somatuline Depot [lanreotide injection], dopamine agonist [e.g., cabergoline, bromocriptine], or Somavert [pegvisomant injection]). Reference ranges for IGF-1 vary among laboratories.
 - C) The medication is prescribed by or in consultation with an endocrinologist.
2. **Carcinoid Syndrome.** Approve for 1 year if the medication is prescribed by or in consultation with an oncologist, endocrinologist, or gastroenterologist.
3. **Neuroendocrine Tumor(s) [NETs] of the Gastrointestinal Tract, Lung, Thymus (Carcinoid Tumors), and Pancreas (including glucagonomas, gastrinomas, vasoactive intestinal peptides-secreting tumors [VIPomas], insulinomas).** Approve for 1 year if the medication is prescribed by or in consultation with an oncologist, endocrinologist, or gastroenterologist.

Other Uses with Supportive Evidence

4. **Pheochromocytoma and Paraganglioma.** Approve for 1 year if the medication is prescribed by or in consultation with an endocrinologist, oncologist, or neurologist.
- II. Coverage of lanreotide subcutaneous injection is recommended in those who meet one of the following criteria:

FDA-Approved Indications

1. **Acromegaly.** Approve for 1 year if the patient meets ALL of the following (A, B, and C):
- A) Patient meets ONE of the following (i, ii, or iii):
 - i. Patient has had an inadequate response to surgery and/or radiotherapy; OR
 - ii. Patient is NOT an appropriate candidate for surgery and/or radiotherapy; OR
 - iii. Patient is experiencing negative effects due to tumor size (e.g., optic nerve compression); AND
 - B) Patient has (or had) a pre-treatment (baseline) insulin-like growth factor-1 (IGF-1) level above the upper limit of normal (ULN) based on age and gender for the reporting laboratory; AND
Note: Pre-treatment (baseline) refers to the IGF-1 level prior to the initiation of any somatostatin analog (e.g., Mycapssa [octreotide delayed-release capsules], an octreotide acetate injection product [e.g., Bynfezia Pen, Sandostatin {generics}, Sandostatin LAR Depot], Signifor LAR [pasireotide injection], Somatuline Depot [lanreotide injection], dopamine agonist [e.g., cabergoline, bromocriptine], or Somavert [pegvisomant injection]). Reference ranges for IGF-1 vary among laboratories.
 - C) The medication is prescribed by or in consultation with an endocrinologist.

Neuroendocrine Tumor(s) [NETs] of the Gastrointestinal Tract, Lung, Thymus (Carcinoid Tumors), and Pancreas (including glucagonomas, gastrinomas, vasoactive intestinal peptides-secreting tumors [VIPomas], insulinomas). Approve for 1 year if the medication is prescribed by or in consultation with an oncologist, endocrinologist, or gastroenterologist.

CONDITIONS NOT RECOMMENDED FOR APPROVAL

Coverage of lanreotide products is not recommended in the following situations:

1. Coverage is not recommended for circumstances not listed in the Recommended Authorization Criteria. Criteria will be updated as new published data are available.

REFERENCES

1. Somatuline[®] Depot subcutaneous injection [prescribing information]. Basking Ridge, NJ: Ipsen; February 2023.
2. Lanreotide subcutaneous injection [prescribing information]. Warren, NJ: Cipla; September 2023.
3. The NCCN Neuroendocrine and Adrenal Tumors Clinical Practice Guidelines in Oncology (version 1.2023 – August 2, 2023). © 2023 National Comprehensive Cancer Network. Available at: <http://www.nccn.org>. Accessed May 10, 2024.